

Magnesium Complex



**MUSCLE
DUMMIES**

Clinical Applications

- Supports Healthy Brain Magnesium Levels*
- Supports Healthy Synapse Number and Function*
- Supports Cognitive Health*
- Supports Stress Management, Sleep Quality, and a Healthy Mood*
- Helps Ensure an Optimal Magnesium Intake for Overall Health*

*Magnesium Complex features magnesium L-threonate (as Magtein®), the only form of magnesium known to cross the blood-brain barrier. Complemented by highly absorbable Albion® di-magnesium malate and TRAACS® magnesium lysinate glycinate chelate, this formula is designed to boost the brain's magnesium level. Magnesium is vital to supporting brain health and promoting healthy nervous system functions.**

All Muscle Dummies LLC Formulas Meet or Exceed cGMP Quality Standards

Discussion

Magnesium has numerous critical roles in human physiology, and its intake is imperative for supporting overall health. It is a cofactor in more than 300 enzymatic reactions that regulate essential functions, including energy production, blood pressure, blood glucose, bone development, and muscle and nerve function. Magnesium is widely available in foods such as whole grains, nuts and seeds, legumes, and green leafy vegetables.¹ In the United States, dietary surveys show magnesium intake is consistently below age-specific minimum recommended levels. An analysis of data from the National Health and Nutrition Examination Survey (NHANES) found that nearly half of Americans ingest less magnesium from food and beverages than is recommended.² Insufficient intake of magnesium is implicated in a wide range of health concerns, including those that affect the brain and nervous system.*¹

Supplemental magnesium can help ensure optimal intake, yet absorption pathways vary among the different forms, and some types can cause gastrointestinal (GI) distress in those with impaired digestion. Factors affecting absorption, tolerance, and retention of supplemental magnesium include the status of magnesium stores in the body and the type or form of magnesium ingested.*³

Magtein® (Patented Magnesium L-Threonate)

More than 10 years of research at MIT went into developing Magtein, a magnesium compound comprised of magnesium and threonic acid commonly known as magnesium L-threonate. Magnesium L-threonate has been suggested to be more bioavailable than other sources of magnesium and has been studied for its unique ability to cross the blood-brain barrier delivering magnesium to brain cells.*⁴⁻⁶

Magnesium Lysinate Glycinate Chelate

Amino acids are used as chelating agents for magnesium to enhance bioavailability, stability, and safety. Chelates increase mineral bioavailability, protect magnesium from being bound in the GI tract, and lower the risk for osmotic diarrhea, which can occur in some forms.³

As an example of the increased bioavailability and tolerability of chelated minerals, the effect of magnesium diglycinate—a chelated form of magnesium bound to 2 glycine molecules—was assessed in subjects (N = 12) with ileal resections. Data from the results suggested that this form of chelated magnesium is likely absorbed intact via a dipeptide transport pathway in different areas of the gut than magnesium oxide, rendering it potentially more bioavailable in addition to being more suitable for those with intestinal resections.⁷ Another study evaluating magnesium absorbability in healthy human subjects (N = 50) demonstrated that magnesium amino acid chelate had better short-term (24 hours) and long-term (60 days) absorbability when compared with magnesium oxide, but the results were not statistically significant.*⁸

Magnesium lysinate glycinate chelate is a mineral amino acid chelate in which magnesium is bound to the amino acids lysine and glycine to create a complex that is readily absorbed across the intestinal wall like other chelated forms.*

Di-Magnesium Malate

This form of magnesium contains malic acid, which also forms a complex with magnesium. In a scientific opinion paper, the European Food Safety Authority (EFSA) cited an unpublished, randomized trial in healthy volunteers (N = 14) in which the bioavailability and tolerability of 150 mg of magnesium from bisglycinate chelate, buffered bisglycinate chelate, and di-magnesium malate were found to be significantly higher than magnesium oxide.⁹ In an animal study, 5 magnesium preparations were evaluated by examining time-dependent absorption, tissue penetration, and effect on the behavior of the animals. Pharmacokinetically, the area under the curve calculation was highest for magnesium malate, suggesting enhanced bioavailability. Further research is needed in humans to assess the dose and effects of di-magnesium bound to malate on specific tissues and behavior.*¹⁰

Brain Health and Cognitive Function

Magnesium has been suggested to play a role in promoting synaptic plasticity in the brain, helping brain cells respond to signals with clarity and vigor without being overactivated. Cognitive decline has been linked to loss of synapse functionality*.¹¹

Ingesting conventional magnesium compounds does not appear to elevate brain magnesium. However, animal studies have shown that magnesium L-threonate crosses the blood-brain barrier, resulting in increased magnesium in extracellular fluid and deposits in neural synapses, increased neural synaptic density, and improved cognitive function.^{4,12-14} Results from a study in rodents suggested that L-threonate is the only ligand to efficiently transport magnesium into cerebrospinal fluid and then into neurons.⁶ To gain a deeper understanding of the mechanisms involved, animal research also found that magnesium L-threonate increased receptor signaling, specifically, the signaling of the NR2B-containing N-methyl-D-aspartate (NMDA) receptor. These NMDA receptors are rich in the hippocampus and play a pivotal role in memory processes.⁴ Several preclinical animal studies that used assessments, such as the novel object recognition test (NORT), T-maze, Morris water maze, conditioned fear memory, and conditioned taste aversion, have validated the effectiveness of magnesium L-threonate in animals. In these studies, researchers demonstrated that when brain magnesium levels were increased, significant benefits were detected in multiple aspects of learning and memory in young and aged rodents.*¹³⁻¹⁵

Continued on next page

***These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure, or prevent any disease.**

Distributed By: Muscle Dummies LLC
228 Park Ave S #702394,
New York, New York 10003
info@muscledummies.com

Magnesium Complex



Supplement Facts

Serving Size: 3 Capsules
Servings Per Container: 30

	Amount Per Serving	%Daily Value
Magnesium (as di-magnesium malate ^{S1} , magnesium L-threonate ^{S2} , and magnesium lysinate glycinate chelate ^{S1})	200 mg	48%
Magnesium L-threonate ^{S2}	1 g	**

** Daily Value not established.

Other Ingredients: Capsule (hypromellose and water), hydroxypropyl cellulose, ascorbyl palmitate, and silica.

S1. Albion® and TRAACS® are registered trademarks of Albion Laboratories, Inc.

S2. Magtein® is protected under a family of U.S. Patents and Pending Patents and is protected worldwide. Magtein® is a registered trademark of Magceutics®, Inc. and is distributed exclusively by AIDP, Inc.



Directions

Take three capsules once or twice daily, or as directed by your healthcare professional.

Consult your healthcare professional before use. Individuals taking medication should discuss potential interactions with their healthcare professional. Do not use if tamper seal is damaged.

Formulated To Exclude:

Wheat, gluten, corn, yeast, soy, animal and dairy products, fish, shellfish, peanuts, tree nuts, egg, sesame, ingredients derived from genetically modified organisms (GMOs), artificial colors, artificial sweeteners, and artificial preservatives.

References

1. Magnesium fact sheet. National Institutes of Health. Updated June 2, 2022. Accessed October 13, 2023. <https://ods.od.nih.gov/factsheets/Magnesium-HealthProfessional/>
2. Usual nutrient intake from food and beverages, by gender and age, what we eat in America, NHANES 2013-2016. USDA, Agricultural Research Service. 2019. Accessed https://www.ars.usda.gov/ARSUserFiles/80400530/pdf/usual/Usual_Intake_gender_WWEIA_2013_2016.pdf
3. Siebrecht S. *Int J Orthomol Relat Med* 2013;144:1-6.
4. Slutsky I, Abumaria N, Wu LJ, et al. *Neuron*. 2010;65(2):165-177. doi:10.1016/j.neuron.2009.12.026
5. Zhang C, Hu Q, Li S, et al. *Nutrients*. 2022;14(24):5235. doi:10.3390/nu14245235
6. Sun Q, Weinger JG, Mao F, et al. *Neuropharmacology*. 2016;108:426-439. doi:10.1016/j.neuropharm.2016.05.006
7. Schuette SA, Lashner BA, Janghorbani M. *JPEN J Parenter Enteral Nutr*. 1994;18(5):430-435. doi:10.1177/0148607194018005430
8. Walker AF, Marakis G, Christie S, et al. *Magnes Res*. 2003;16(3):183-191.
9. EFSA Panel on Food Additives and Nutrient Sources added to Food (ANS); Younes M, Aggett P, et al. *EFSA J*. 2018;16(6):e05292. doi:10.2903/j.efsa.2018.5292
10. Uysal N, Kizildag S, Yuca Z, et al. *Biol Trace Elem Res*. 2019;187(1):128-136. doi:10.1007/s12011-018-1351-9
11. András E, Páli N, Molnár Z, et al. *J Alzheimers Dis*. 2005;7(4):273-284. doi:10.3233/jad-2005-7402
12. Slutsky I, Sadeghpour S, Li B, et al. *Neuron*. 2004;44(5):835-849. doi:10.1016/j.neuron.2004.11.013
13. Li W, Yu J, Liu Y, et al. *Mol Brain*. 2014;7:65. doi:10.1186/s13041-014-0065-y
14. Abumaria N, Yin B, Zhang L, et al. *J Neurosci*. 2011;31(42):14871-14881. doi:10.1523/JNEUROSCI.3782-11.2011
15. Mickley GA, Hoxha N, Luchsinger JL, et al. *Pharmacol Biochem Behav*. 2013;106:16-26. doi:10.1016/j.pbb.2013.02.019
16. Liu G, Weinger JG, Lu ZL, et al. *J Alzheimers Dis*. 2016;49(4):971-990. doi:10.3233/JAD-150538
17. Eby GA, Eby KL. *Med Hypotheses*. 2006;67(2):362-370. doi:10.1016/j.mehy.2006.01.047
18. Pochwat B, Szweczyk B, Sowa-Kucma M, et al. *Int J Neuropsychopharmacol*. 2014;17(3):393-405. doi:10.1017/S1461145713001089
19. Abumaria N, Luo L, Ahn M, et al. *Behav Pharmacol*. 2013;24(4):255-263. doi:10.1097/FBP.0b013e32836357c7
20. Held K, Antonijevic IA, Künzel H, et al. *Pharmacopsychiatry*. 2002 Jul;35(4):135-143. doi:10.1055/s-2002-33195
21. Abbasi B, Kimiagar M, Sadeghnia K, et al. *J Res Med Sci*. 2012;17(12):1161-1169
22. Hornyak M, Voderholzer U, Hohagen F, et al. *Sleep*. 1998;21(5):501-505. doi:10.1093/sleep/21.5.501
23. Nielsen FH, Johnson LK, Zeng H. *Magnes Res*. 2010;23(4):158-68. doi:10.1684/mrh.2010.0220
24. Saba S, Faizi F, Sepandi M, et al. *Magnes Res*. 2022;35(2):62-70. doi:10.1684/mrh.2022.0503

***These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure, or prevent any disease.**

Distributed By: Muscle Dummies LLC
228 Park Ave S #702394,
New York, New York 10003
info@muscleddummies.com

DRS-356
REV. 02/26/24